



Contents lists available at SciVerse ScienceDirect

International Journal of Psychophysiology

journal homepage: www.elsevier.com/locate/ijpsycho

Tracking EEG changes in response to alpha and beta binaural beats

D. Vernon ^{a,*}, G. Peryer ^b, J. Louch ^a, M. Shaw ^a^a Department of Applied Social Sciences, Canterbury Christ Church University, Canterbury, Kent CT1 1QU, UK^b Faculty of Medicine and Health Sciences, University of East Anglia, Norwich Research Park, Norwich NR4 7TJ, UK

ARTICLE INFO

Article history:

Received 15 June 2012

Received in revised form 5 October 2012

Accepted 11 October 2012

Available online xxxxx

Keywords:

Binaural beat

EEG

Alpha

Beta

Entrainment

ABSTRACT

A binaural beat can be produced by presenting two tones of a differing frequency, one to each ear. Such auditory stimulation has been suggested to influence behaviour and cognition via the process of cortical entrainment. However, research so far has only shown the frequency following responses in the traditional EEG frequency ranges of delta, theta and gamma. Hence a primary aim of this research was to ascertain whether it would be possible to produce clear changes in the EEG in either the alpha or beta frequency ranges. Such changes, if possible, would have a number of important implications as well as potential applications. A secondary goal was to track any observable changes in the EEG throughout the entrainment epoch to gain some insight into the nature of the entrainment effects on any changes in an effort to identify more effective entrainment regimes. Twenty two healthy participants were recruited and randomly allocated to one of two groups, each of which was exposed to a distinct binaural beat frequency for ten 1-minute epochs. The first group listened to an alpha binaural beat of 10 Hz and the second to a beta binaural beat of 20 Hz. EEG was recorded from the left and right temporal regions during pre-exposure baselines, stimulus exposure epochs and post-exposure baselines. Analysis of changes in broad-band and narrow-band amplitudes, and frequency showed no effect of binaural beat frequency eliciting a frequency following effects in the EEG. Possible mediating factors are discussed and a number of recommendations are made regarding future studies, exploring entrainment effects from a binaural beat presentation.

© 2012 Published by Elsevier B.V.

1. Introduction

A binaural beat occurs when two carrier tones of distinct frequency are presented separately, one to each ear, and the resulting perception is of a single tone with a frequency midway between the two carrier tones that waxes and wanes in amplitudes at a rate equal to the difference between them (Oster, 1973). For instance, if a clear tone of 400 Hz is presented to the left ear and a tone of 410 Hz is presented to the right ear this produces a perceived frequency of 405 Hz which modulates in amplitude with a frequency of 10 Hz. This modulated signal is often referred to as a beat and can be produced when the difference in frequency between the two carrier tones ranges from between 2 and 30 Hz (Perrott and Nelson, 1969). Furthermore, there are suggestions that carrier tones of a frequency ranging from 200 to 900 Hz may be more effective at eliciting binaural beats than carrier tones that exceed 1 kHz (Wahbeh et al., 2007; Pratt et al., 2010). Nevertheless, it has recently been suggested that the perceptual strength of a binaural beat signal remains relatively weak (Grose et al., 2012).

A number of reports have suggested that listening to binaural beats can influence behaviour and cognition in a variety of ways. For instance, reports show that exposure to binaural beats has led to

improved vigilance (Lane et al., 1998) and memory (Kennerly, 1996), increased levels of hypnotic susceptibility (Brady and Stevens, 2000) and led to reductions in self-reports of anxiety (Le Scouarnec et al., 2001; Padmanabhan et al., 2005) and severity ratings of tinnitus (David et al., 2010). However, others report that exposure to binaural beats has failed to influence hypnotic susceptibility (Stevens et al., 2003), produced no change in blood pressure or heart rate (Carter, 2008) and did not reduce the symptoms of children diagnosed with attention deficit hyperactivity disorder (Kennel et al., 2010). It is difficult to know why such different outcomes emerge and this in part highlights a more general lack of research regarding the possible effects of listening to binaural beats. In part, such differences may occur due to the distinct methodological approaches adopted which in turn may influence the entrainment potential of binaural beats. This is further limited by the fact that only two of the above studies (Brady and Stevens, 2000; Stevens et al., 2003) monitored for potential changes in electroencephalographic (EEG) activity as a function of exposure to binaural beats.

A central assumption of binaural beats is that they can elicit an entrainment effect, sometimes referred to as a frequency following response (FFR), in the electrocortical activity of the brain and thereby represent a mechanism for potential behaviour change (for a review see Vernon, 2009). According to Hink et al. (1980) the FFR arises from the brain stem. They suggest that the FFR stems from converging input from overlapping neuronal populations and that this can be

* Corresponding author.

E-mail address: david.vernon@canterbury.ac.uk (D. Vernon).

directly influenced by binaural beats. It is not clear as yet precisely how widespread the changes in electrocortical activity may be following exposure to binaural beats as there are a number of inconsistencies in the literature. For example, some have reported changes in amplitude at the central midline region (Cz) following exposure to binaural beats (Schwarz and Taylor, 2005) whilst others have failed to find any evidence of such a change (Wahbeh et al., 2007). These inconsistencies may stem from differences in methodology, in particular the nature and frequency of the carrier tones used to present the binaural beats as well as the duration of exposure. Nevertheless, there is an emerging consensus that exposure to binaural beats may lead to changes in brain activity in the temporal regions of the brain (Karino et al., 2004, 2006; Pratt et al., 2010). For instance, Karino et al. (2006) reported dominant activation of both left and right temporal areas, which contain the auditory cortex, following exposure to binaural beats in the theta range (4 Hz and 6.6 Hz). More recently Pratt et al. (2010) noted that the source of the oscillations elicited by exposure to binaural delta (3 Hz) and theta (6 Hz) beats were predominantly located to the left temporal region. Such findings are consistent with early suggestions that the temporal lobe is electrically more labile than other regions of the cortex (Bear, 1979). Hence, we have focused on possible changes in the EEG at the left and right temporal regions.

Despite such encouraging findings, to date changes have only been reported in three of the main traditional EEG frequency ranges. This includes changes in delta (Kennerly, 1996; Pratt et al., 2010), theta (Brady and Stevens, 2000; Karino et al., 2004, 2006; Pratt et al., 2010) and gamma (Schwarz and Taylor, 2005). However, no changes have yet been reported in the key EEG frequencies of alpha and beta. Hence, the primary aim of this study was to examine the effects of alpha and beta binaural beats on the EEG. By doing this we hoped to ascertain whether such a technique has the potential to increase activity in specific targeted components of the EEG which would have a number of practical applications with regard to performance enhancement, as well as clinical treatment (for a review see Vernon, 2009).

The possible entrainment and modification of alpha and beta EEG activities would present with some interesting possibilities. For instance, there is a growing body of research showing that alpha activity, far from representing cortical idling, may in fact provide an index of neural efficiency (see Vernon, 2009). This idea is supported by research showing that individuals classified as more intelligent exhibit greater levels of alpha power (Anokhin and Vogel, 1996; Jausovec, 1996; Doppelmayr et al., 2005) and that induced increases in alpha power using EEG biofeedback (Hanslmayr et al., 2005) and repetitive transcranial magnetic stimulation (Klimesch et al., 2003) have been associated with improved attentional processing. In addition, beta EEG activity has been implicated in attentional processing, much of which is based on attempts to normalise the EEG of those with attentional dysfunction (see e.g., Lubar et al., 1995a, 1995b). Support for this comes from research focusing on enhancing beta power using EEG biofeedback which has been shown to improve attentional processing in both clinical populations (Fuchs et al., 2003; Monastra et al., 2006) and healthy participants (Rasey et al., 1996; Egner and Gruzelie, 2004). In addition, research has shown increases in beta (12–30 Hz) amplitude when participants attend to a stimulus and plan a response (Saleh et al., 2010). These findings would support the use of binaural beats as a plausible mechanism for enhancing EEG power within a targeted frequency in an attempt to normalise and/or enhance cognition, if exposure to such sounds elicited clear FFR's in the EEG.

A secondary goal of the current research was to track the changes in the EEG throughout the period of exposure to the binaural beats in an attempt to obtain some insight into the possible nature of the brain's response to such sounds. Such information could prove helpful in designing and refining future entrainment protocols. It is not

clear as yet if and how, the brain will respond to such audio stimulation. For instance, previous research is inconsistent with some showing that theta (6.6 Hz) binaural beats are capable of eliciting changes when measured using a very short exposure window of 1000 ms (Karino et al., 2004), whilst others have failed to find any evidence of cortical entrainment when theta (7 Hz) binaural beats were constantly presented for 2 min (Goodin et al., 2012), or for periods of up to 30 min (Wahbeh et al., 2007). It may be that continuous and prolonged exposure to a binaural beat leads the brain to habituate to the signal, reducing the possibility of identifying an FFR as time progresses. Such a possibility would be consistent with research showing habituation of evoked potentials to repeating auditory stimuli (e.g., Barry et al., 1992). In an attempt to address this we devised a novel exposure procedure which involved presenting the binaural beat signal in ten 1-minute segments interleaved with a single carrier tone set at the same amplitude and frequency as the perceived frequency of the binaural beats. Given that the alpha binaural beat was created using carrier tones of 395 and 405 Hz, producing a perceived frequency of 400 Hz modulating at 10 Hz, this meant that each 1-minute segment of binaural beats was interleaved with a single pure tone played at 400 Hz. Tones centred around 400 Hz were used as Oster (1973) suggested that binaural beats may be better perceived with the carrier tones being close to 400 Hz (see also Goodin et al., 2012). Such a procedure would not only counteract the possible habituation effects of continuous and prolonged exposure to the binaural beats but would allow us to track any possible changes in the EEG over time.

2. Method

2.1. Participants

Twenty two participants were recruited from the Psychology undergraduate cohort at Canterbury Christ Church University. Participants' ages ranged from 18 to 32 (mean 20 y) with 13 females and 9 males. Participants were randomly allocated to either the *alpha-entrainment* group or the *beta-entrainment* group,¹ however, due to timing issues this resulted in 12 participants in the *alpha-entrainment* group and 10 in the *beta-entrainment* group. All participants had normal or corrected to normal vision, no reported hearing deficits/difficulties and were screened to exclude individuals with a history/family history of epilepsy, psychological problems and those taking prescribed medication. Participants received course credits for completing the study.

2.2. Binaural beats

Previous research has suggested that binaural beats may be perceived with greater clarity when the carrier tones are higher than 100 Hz (Perrott and Nelson, 1969) and closer to 400 Hz (Oster, 1973). Hence, two binaural beat recordings were created using Adobe Audition. The *alpha* entrainment recording consisted of ten 1-minute segments containing two sinusoidal tones, one at 395 Hz and one at 405 Hz to produce a 10 Hz binaural beat, which was interspersed ten 1-minute segments of a single carrier tone at 400 Hz. The *beta* entrainment recording consisted of ten 1-minute segments containing two sinusoidal tones, one at 390 Hz and one at 410 Hz to produce a 20 Hz binaural beat, which was interspersed with ten 1-minute segments of a single carrier tone at 400 Hz. All sound segments contained a 10 ms attack and decay period.

¹ Participants were exposed to a single binaural beat frequency to avoid potential confounding carry over effects whereby entrainment at one frequency influences the EEG outside of that frequency range (see e.g., Egner and Gruzelie, 2003; Lavallee et al., 2011).

208 2.3. EEG acquisition

209 EEG signals were acquired using a Nexus-10 DC-coupled EEG
 210 portable amplifier incorporating a 24-bit A/D converter with BioTrace
 211 software (MindMedia, Netherlands). EEG recording was sampled at
 212 256 Hz using two monopolar/referential sensors, one attached to T3
 Q13213 (according to international 10/20 system: Jasper, 1958) and referenced
 214 to the left mastoid and the other attached to T4 and referenced to the
 215 right mastoid, with a ground sensor which was placed at Fz. Targeted
 216 scalp regions were prepared using NuPrep abrasive gel and sensors
 217 attached using Ten20 electrode conducting paste. All sampled data
 218 were initially subject to off-line voltage threshold artefacting to help re-
 219 move possible interference from ocular, head and muscle movements.
 220 EEG data analysis was conducted off-line with the raw trace digitally
 221 filtered using an IIR Bandpass Butterworth (3rd order) filter to extract
 222 broadband alpha (8–12 Hz), narrow-band alpha (9.5–10.5 Hz), broad-
 223 band beta (18–22 Hz) and narrow-band beta (18.5–20.5 Hz) mean
 224 amplitude exported at 32 samples/s. Peak frequency was determined
 225 using a fast four transform to identify the dominant frequency using
 226 4 second epochs within a range of 8–12 Hz for alpha and 18–22 Hz
 227 for beta. Means were then computed for each of the target EEG compo-
 228 nents of interest for each minute of the recording.

229 2.4. Procedure

230 Participants were informed of the purpose of the study and were
 231 randomly allocated to one of the two auditory conditions (alpha,
 232 beta). They then completed each of the three stages whilst remaining
 233 seated. In the initial stage a 2-minute eyes open resting EEG baseline
 234 and a 2-minute eyes closed EEG resting baseline was recorded, with
 235 order counterbalanced across participants. In stage 2 participants
 236 were shown a nature DVD with the sound muted whilst simultaneously
 237 presented with a binaural beat recording via on-ear headphones.
 238 Each binaural beat recording consisted of ten 1-minute segments of bin-
 239 aural beats interspersed with ten 1-minute segments of a single carrier
 240 tone. The order of these was counterbalanced across participants.
 241 Throughout the auditory exposure period participants' EEG was
 242 recorded from T3 and T4. During the final stage a 2-minute eyes open
 243 and 2-minute eyes closed resting EEG baselines were again recorded.

244 3. Results

245 Mean amplitude and frequency measures are reported separately
 246 for each group and for baseline and entrainment measures. For all
 247 analyses a Greenhouse–Geisser correction was used if Mauchley's
 248 test of sphericity was significant and corrected df are reported to
 249 two decimal places.

250 3.1. Alpha EEG

251 A 2 (Time: *Pre* vs. *Post*) × 2 (Hemisphere: *Left* vs. *Right*) repeated
 252 measures analysis of variance (ANOVA) was conducted on mean broad-
 253 band (8–12 Hz) amplitude, mean narrow-band (9.5 Hz–10.5 Hz)
 254 amplitude and peak alpha (8–12 Hz) frequency separately for eyes
 255 open and eyes closed resting baselines.

256 3.2. Eyes open

257 For broadband amplitude there was a marginal effect of Time, F
 258 (1,11) = 4.394, $p = 0.06$, $Mse = 17.23$, $\eta^2 = 0.285$, showing a drop in
 259 amplitude from the pre to post entrainment from 8.58 μV to 7.34 μV .
 260 For both narrow-band amplitude and peak frequency there were no
 261 main or interaction effects (all $ps > 0.18$).

3.3. Eyes closed

262 For both broadband and narrow-band amplitudes there were no
 263 main or interaction effects (all $ps > 0.14$). For peak alpha frequency
 264 there was a main effect of the Hemisphere, $F(1,11) = 5.065$, $p < 0.05$,
 265 $Mse = 0.083$, $\eta^2 = 0.315$, showing a higher peak alpha frequency in
 266 the left hemisphere (9.81 Hz) compared to the right (9.72 Hz). There
 267 were no other main or interaction effects (all $ps > 0.45$).
 268

3.4. Alpha entrainment

269 A 2 (Hemisphere: *Left* vs. *Right*) × 10 (Time: 1 min to 10 min) × 2
 270 (Signal: *BB-on* vs. *BB-off*) repeated measures analysis of variance
 271 (ANOVA) was conducted on the mean broadband (8–12 Hz) amplitude,
 272 mean narrow-band (9.5 Hz–10.5 Hz) amplitude and peak alpha
 273 (8–12 Hz) frequency separately. Descriptives for each of these variables
 274 can be seen below in Table 1.
 275

276 There were no main or interaction effects for broadband ampli-
 277 tude (all $ps > .107$), narrow-band amplitude (all $ps > .112$) or peak
 278 alpha frequency (all $ps > .027$).
 279

280 Given the failure to find any effects of entrainment we re-examined
 281 the initial eyes-open resting baseline alpha amplitudes and conducted a
 282 median split to identify those with low (<8.5 μV) resting baselines as
 283 research focusing on audio-visual entrainment has suggested that
 284 only those with low baseline alpha activity may exhibit entrainment
 285 effects in the EEG (Rosenfeld et al., 1997).²

286 A re-analysis utilising a 2 (Signal: *BB-on* vs. *BB-off*) × 10 (Time:
 287 1 min to 10 min) repeated measures analysis of variance (ANOVA)
 288 was conducted on mean broadband (8–12 Hz) amplitude for those
 289 with a low resting baseline level. This showed a main effect of Signal,
 290 $F(1,4) = 13.952$, $p < 0.05$, $Mse = 0.203$, $\eta^2 = 0.777$, showing greater
 291 alpha amplitude during the *BB-off* phase compared to the *BB-on*
 292 phase (4.84 μV and 4.75 μV respectively).³ There were no other
 293 main or interaction effects (all $ps > 0.23$).⁴

3.5. Beta EEG

294 A 2 (Time: *Pre* vs. *Post*) × 2 (Hemisphere: *Left* vs. *Right*) repeated
 295 measures analysis of variance (ANOVA) was conducted on mean
 296 broadband (18–20 Hz) amplitude, mean narrow-band (19.5 Hz–
 297 20.5 Hz) amplitude and peak beta (18–20 Hz) frequency separately
 298 for eyes open and eyes closed resting baselines.
 299

3.6. Eyes open

300 For both the broadband and narrow-band amplitudes there was a
 301 Time by Hemisphere interaction. Broadband: $F(1,9) = 22.224$, $p < 0.01$,
 302 $Mse = 13.47$, $\eta^2 = 0.712$; Narrow-band: $F(1,9) = 6.776$, $p < 0.05$, $Mse =$
 303 3.091 , $\eta^2 = 0.43$. Post hoc comparisons using a Bonferroni correction
 304 (0.05/2 = 0.025) examined the changes in amplitude from the *Pre* to
 305 *Post* for the *Left* and *Right* hemispheres separately. These showed a mar-
 306 ginal increase in amplitude from the *Pre* to *Post* for the *Left* hemisphere
 307 (broadband: 7.69 μV and 9.65 μV $p = 0.041$; narrow-band 3.54 μV and
 308 4.56 μV $p = 0.032$) but not the *Right* (broadband: 7.56 μV and 7.20 μV
 309 $p = 0.65$; narrow-band: 3.74 μV and 3.65 μV $p = 0.87$). There were no

² Given the lack of any effects of the hemisphere we collapsed the data across right and left hemispheres and focused only on the eyes open pre entrainment baselines as the entrainment was also conducted with eyes open. Re-analysis of the entrainment effects was also conducted collapsing across hemispheres given the initial null effects and also to reduce variability given the smaller N.

³ A corresponding analysis on those with high resting baselines showed no evidence of any effect of Signal ($p = 0.81$).

⁴ A similar analysis conducted on narrow-band amplitude and frequency revealed no main or interaction effects (all $ps > 0.25$).

Table 1
Means (SD) for alpha broadband (8–12 Hz), narrow-band (9.5–10.5 Hz) amplitude and peak frequency (8–12 Hz) for the ten 1-minute binaural-beat on signal entrainment periods and the ten 1-minute binaural-beat off carrier tone baseline periods.

Time periods				1	2	3	4	5	6	7	8	9	10
Broadband amplitude μV	Right hemisphere	BB-on	6.80 (2.65)	6.56 (2.83)	6.67 (3.23)	6.34 (2.84)	6.08 (2.56)	5.66 (2.43)	6.75 (3.45)	5.91 (3.11)	5.85 (2.44)	6.46 (3.12)	
	Left hemisphere	BB-off	6.89 (2.26)	6.95 (2.45)	6.24 (2.77)	6.41 (3.19)	6.37 (3.18)	6.57 (2.22)	6.31 (3.07)	6.16 (3.46)	5.53 (1.57)	6.06 (3.05)	
	Right hemisphere	BB-on	7.54 (2.87)	6.61 (1.18)	6.92 (2.39)	6.29 (2.00)	6.56 (1.85)	6.30 (1.87)	7.52 (2.35)	6.56 (2.05)	6.24 (1.73)	6.32 (1.41)	
	Left hemisphere	BB-off	7.54 (2.84)	7.36 (3.02)	6.19 (1.55)	6.62 (2.29)	6.49 (1.70)	6.75 (1.95)	6.72 (1.83)	6.85 (1.84)	6.37 (1.64)	6.73 (1.78)	
Narrow-band amplitude μV	Right hemisphere	BB-on	3.39 (1.75)	3.05 (1.44)	3.32 (1.93)	3.08 (1.54)	2.91 (1.32)	2.81 (1.45)	3.40 (2.28)	2.81 (1.55)	2.86 (1.34)	3.16 (1.77)	
	Left hemisphere	BB-off	3.20 (1.11)	3.40 (1.39)	3.10 (1.65)	3.15 (1.79)	3.04 (1.82)	3.16 (1.15)	2.99 (1.52)	2.91 (1.85)	2.56 (0.66)	2.98 (1.79)	
	Right hemisphere	BB-on	3.46 (1.55)	3.01 (0.57)	3.24 (1.21)	2.97 (0.97)	3.01 (0.83)	3.00 (0.95)	3.85 (1.78)	3.05 (1.02)	3.01 (0.96)	3.08 (0.84)	
	Left hemisphere	BB-off	3.67 (1.74)	3.52 (1.79)	2.94 (0.85)	3.03 (1.09)	3.06 (0.90)	3.16 (0.86)	3.18 (1.01)	3.45 (1.05)	3.05 (0.75)	3.22 (0.87)	
Peak frequency Hz	Right hemisphere	BB-on	10.0 (0.26)	9.96 (0.29)	9.84 (0.54)	9.89 (0.44)	9.85 (0.59)	10.0 (0.39)	9.94 (0.51)	9.92 (0.49)	9.97 (0.52)	9.95 (0.32)	
	Left hemisphere	BB-off	10.0 (0.28)	9.96 (0.28)	9.93 (0.46)	9.97 (0.36)	9.84 (0.47)	10.0 (0.48)	9.91 (0.49)	9.92 (0.59)	10.0 (0.35)	9.98 (0.49)	
	Right hemisphere	BB-on	10.0 (0.39)	10.0 (0.50)	10.0 (0.61)	9.94 (0.55)	9.97 (0.47)	9.93 (0.46)	10.0 (0.33)	9.96 (0.41)	10.1 (0.52)	9.99 (0.51)	
	Left hemisphere	BB-off	10.0 (0.37)	10.1 (0.44)	10.0 (0.52)	9.90 (0.57)	9.96 (0.44)	10.0 (0.42)	10.1 (0.42)	10.1 (0.43)	9.90 (0.28)	9.97 (0.44)	

other main effects for the amplitude (all $p > 0.20$) and no main or interaction effects for the peak beta frequency (all $p > 0.54$).

3.7. Eyes closed

For both broadband and narrow-band amplitudes there were main effects of Time. Broadband: $F(1,9) = 20.73$, $p < 0.001$, $Mse = 34.93$, $\eta^2 = 0.70$; narrow-band: $F(1,9) = 24.33$, $p < 0.001$, $Mse = 15.34$, $\eta^2 = 0.73$, showing a drop in amplitudes from the pre to post entrainment (broadband: from 8.72 μV to 6.85 μV ; narrow-band from 4.65 μV to 3.41 μV). For the narrow-band amplitude only there was also a marginal effect of the Hemisphere, $F(1,9) = 4.69$, $p = 0.058$, $Mse = 11.97$, $\eta^2 = 0.34$, with less amplitude in the *Left* hemisphere compared to the *Right* (4.58 μV to 3.48 μV respectively). There were no other main or interaction effects for amplitude (all $p > 0.11$) and no main or interaction effects for the peak beta frequency (all $p > 0.18$).

3.8. Beta entrainment

A 2 (Hemisphere: *Left* vs. *Right*) \times 10 (Time: 1 min to 10 min) \times 2 (Signal: *BB-on* vs. *BB-off*) repeated measures analysis of variance (ANOVA) was conducted on the mean broadband (18–22 Hz) amplitude, mean narrow-band (19.5 Hz–20.5 Hz) amplitude and peak beta (18–22 Hz) frequency separately. Descriptives for each of these variables can be seen below in Table 2.

For both broadband and narrow-band amplitudes there was a marginal effect of Time: broadband, $F(9,81) = 2.675$, $p = 0.082$, $Mse = 15.88$, $\eta^2 = 0.23$; narrow-band, $F(2.62, 23.65) = 2.879$, $p = 0.063$, $Mse = 5.50$, $\eta^2 = 0.24$, showing a drop in amplitude from *time-1* to *time-10* (broadband: 7.52 μV to 5.65 μV ; narrow-band 3.41 μV to 2.19 μV). There was also a marginal effect of the Hemisphere:

broadband, $F(1,9) = 4.654$, $p = 0.059$, $Mse = 159.51$, $\eta^2 = 0.34$; narrow-band, $F(1,9) = 4.152$, $p = 0.072$, $Mse = 51.77$, $\eta^2 = 0.31$, showing greater amplitude in the *Left* hemisphere compared to the *Right* (broadband: 6.55 μV to 5.29 μV ; narrow-band 2.90 μV to 2.18 μV). For the narrow-band amplitude only there was also a significant Time by Hemisphere interaction, $F(9,81) = 2.149$, $p < 0.05$, $Mse = 0.83$, $\eta^2 = 0.19$ [broadband interaction was $p = 0.131$]. This was explored by conducting a 10 (Time: 1 min to 10 min) \times 2 (Signal: *BB-on* vs. *BB-off*) repeated measures analysis of variance (ANOVA) on the mean narrow-band (19.5 Hz–20.5 Hz) amplitude for the *Left* and *Right* hemispheres separately. For the *Left* hemisphere this revealed a main effect of Time, $F(2.35, 21.15) = 3.684$, $p < 0.05$, $Mse = 3.95$, $\eta^2 = 0.29$ with pairwise comparisons showing a reduction in the amplitude from *time-1* to *time-8* (4.02 μV to 2.55 μV ; $p < 0.05$) and marginal reductions from *time-1* to *time-9* (4.02 μV to 2.79 μV ; $p = 0.087$) and from *time-1* to *time-10* (4.02 μV to 2.48 μV ; $p = 0.095$). There were no main or interaction effects for the *Right* hemisphere (all $p > 0.15$).

For the beta peak frequency there was a significant Time by Hemisphere interaction, $F(9,81) = 2.117$, $p < 0.05$, $Mse = 0.082$, $\eta^2 = 0.19$ but no other main or interaction effects (all $p > 0.13$). The interaction was explored by conducting a 10 (Time: 1 min to 10 min) \times 2 (Signal: *BB-on* vs. *BB-off*) repeated measures analysis of variance (ANOVA) on the mean peak beta frequency for the *Left* and *Right* hemispheres separately. However, this produced no main or interaction effects for either the *Left* hemisphere (all $p > 0.21$) or the *Right* hemisphere (all $p > 0.13$).

4. Discussion

Exposure to ten 1-minute segments of either alpha (10 Hz) or beta (20 Hz) binaural beats produced no clear frequency following

Table 2
Means (SD) for beta broadband (18–22 Hz), narrow-band (19.5–20.5 Hz) amplitude and peak frequency (18–22 Hz) for the ten 1-minute binaural-beat on signal entrainment periods and the ten 1-minute binaural-beat off carrier tone baseline periods.

Time periods				1	2	3	4	5	6	7	8	9	10
Broadband amplitude μV	Right hemisphere	BB-on	7.26 (4.41)	5.64 (3.58)	5.34 (3.08)	5.05 (2.39)	5.08 (2.33)	4.66 (1.84)	4.85 (2.00)	5.20 (2.43)	5.57 (2.53)	4.81 (1.49)	
	Left hemisphere	BB-off	5.75 (2.38)	5.51 (3.36)	5.34 (3.48)	5.20 (2.46)	5.00 (2.20)	4.58 (2.11)	4.83 (2.19)	5.62 (2.91)	5.71 (2.64)	4.71 (1.70)	
	Right hemisphere	BB-on	8.75 (3.28)	7.00 (2.63)	6.37 (2.46)	6.11 (2.11)	6.87 (2.04)	6.61 (1.98)	6.02 (2.41)	5.53 (1.96)	6.30 (3.45)	6.05 (2.61)	
	Left hemisphere	BB-off	8.32 (3.33)	6.79 (2.37)	6.24 (2.56)	6.60 (2.23)	6.70 (2.16)	6.23 (1.79)	6.11 (1.83)	6.23 (3.61)	6.59 (3.27)	5.56 (2.11)	
Narrow-band amplitude μV	Right hemisphere	BB-on	3.09 (1.87)	2.33 (1.29)	2.19 (1.05)	2.17 (1.08)	2.09 (0.85)	1.82 (0.67)	1.94 (0.64)	2.26 (0.96)	2.27 (1.05)	1.92 (0.54)	
	Left hemisphere	BB-off	2.52 (1.05)	2.30 (1.36)	2.20 (1.30)	2.07 (0.84)	1.99 (0.74)	1.81 (0.66)	1.93 (0.95)	2.32 (1.42)	2.45 (1.09)	1.87 (0.53)	
	Right hemisphere	BB-on	4.18 (1.83)	3.12 (1.37)	2.79 (1.20)	2.70 (0.86)	3.09 (1.17)	2.97 (1.18)	2.55 (1.16)	2.47 (0.95)	2.73 (1.74)	2.89 (1.25)	
	Left hemisphere	BB-off	3.86 (1.70)	2.98 (1.13)	2.79 (1.13)	2.91 (1.00)	3.01 (1.02)	2.66 (0.75)	2.73 (0.75)	2.62 (1.78)	2.84 (1.49)	2.36 (0.93)	
Peak frequency Hz	Right hemisphere	BB-on	20.1 (0.46)	20.1 (0.50)	20.3 (0.45)	20.3 (0.44)	20.4 (0.55)	20.5 (0.47)	20.4 (0.58)	20.4 (0.42)	20.4 (0.54)	20.4 (0.66)	
	Left hemisphere	BB-off	20.2 (0.41)	20.3 (0.45)	20.3 (0.60)	20.5 (0.48)	20.5 (0.53)	20.4 (0.51)	20.4 (0.51)	20.3 (0.52)	20.3 (0.47)	20.2 (0.59)	
	Right hemisphere	BB-on	20.2 (0.45)	20.0 (0.60)	20.1 (0.52)	20.1 (0.51)	20.1 (0.53)	20.2 (0.40)	20.0 (0.44)	20.1 (0.46)	20.1 (0.46)	20.1 (0.41)	
	Left hemisphere	BB-off	20.3 (0.42)	20.1 (0.47)	20.0 (0.56)	20.1 (0.53)	20.1 (0.49)	20.0 (0.47)	20.1 (0.43)	20.2 (0.43)	20.2 (0.40)	20.1 (0.43)	

effects in the EEG. For those exposed to the alpha binaural beat there was some evidence of a decrease in resting baseline amplitude from the pre to post entrainment and during the session participants exhibited greater alpha activity during the binaural-beat off phase compared to the on phase. For those exposed to the beta binaural beat there was some evidence of an increase in the resting eyes open baseline beta amplitude for the left hemisphere only. However, during eyes closed resting baseline sessions there was a drop in amplitude from the pre to post entrainment. In addition, during the entrainment session there was a drop in amplitude over time with more robust reductions seen for the narrow-band amplitude in the left hemisphere.

These results provide no clear indication that a frequency following the effect was elicited in the EEG following exposure to either alpha or beta binaural beats. A central assumption of the binaural beats technique is that listening to the sounds elicits an entrainment effect in the EEG and in this way various behaviours may be modified (see Vernon, 2009). However, the results of this study would suggest that exposure to 10 interleaved 1-minute segments of either alpha or beta binaural beats at the perceived frequency of 400 Hz is not sufficient to elicit any such change in the EEG and as such would be expected to have no influence on behaviour. This lack of any clear change in either the alpha or beta EEG frequencies following exposure to binaural beats is consistent with those reporting similar null effects following exposure to the theta (Wahbeh et al., 2007) and alpha (Goodin, et al., 2012) binaural beats. However, this is in contrast to others who have reported changes in the EEG following exposure to the delta/theta (Brady and Stevens, 2000; Karino et al., 2004, 2006; Pratt et al., 2010) and gamma (Schwarz and Taylor, 2005; Grose and Mamo, 2011) frequency ranges. These inconsistent findings suggest that it may be too early to label the technique of binaural beats as completely ineffective as there may be a number of factors that mediate the effectiveness of such an entrainment technique. For instance, it may not be possible for each frequency component of the EEG to be equally entrained, or it may be that participants need to attend to the stimuli for an entrainment effect to occur. In addition, the possible changes elicited by the binaural beats may be more widespread and/or not visible at the temporal regions, or the specific nature and duration of the tones themselves may influence the outcome. Each of these points is discussed in turn below.

It is possible that only certain EEG frequencies exhibit a frequency following effects when exposed to binaural beats. Such a possibility would be consistent with those showing changes in the EEG following exposure to low frequency delta (3 and 4 Hz) and theta (~6 Hz) binaural beats (Karino et al., 2004, 2006; Pratt et al., 2010). In addition, Pratt et al. (2010) reported greater changes in the EEG following exposure to a 3 Hz binaural beat compared to 6 Hz. Such findings could imply that the lower EEG frequencies may be more amenable to change. However, others have reported clear changes in the EEG following exposure to higher gamma (40 Hz) frequency binaural beats (Grose and Mamo, 2011). In addition, combined audio and visual entrainment techniques have been shown to elicit changes in both the alpha (Teplan et al., 2006; Moridis et al., 2010) and beta (Timmerman et al., 1999) frequency ranges of the EEG. Such findings suggest a level of cortical flexibility that is evident across all frequency ranges of the EEG and as such it should be possible to elicit entrainment effects in these frequencies.

Alternatively, it is possible that the effects of binaural beats may be attenuated if the tones are not directly attended to. However, Lane et al. (1998) reported changes in the behaviour following exposure to beta binaural beats when participants remained blind to the presence of the sounds. Unfortunately no EEG measures were taken so it is not clear whether any changes occurred in the EEG. In contrast, Goodin et al. (2012) had participants focus their attention directly on the sounds of the alpha binaural beats and respond when they were able to perceive the beats. Nevertheless, such focused perception

had no effect on the binaural beats which failed to elicit any entrainment effects in the EEG. Interestingly, Schwarz and Taylor (2005) found that when a few of their participants who were musically trained attempted to assign musical intervals to the binaural beat carrier tones by attending to them separately no change was evident in the EEG. However, when the same participants later attended to both carrier tones simultaneously a change in the EEG in response to the binaural beats was evident. This would suggest that individuals, at least those trained in music, may need to 'attend' to the two tones together rather than separately and as such a 'degree' of attention may be helpful. As yet there has been no reported comparison between the effects of attending to binaural beats vs. non-attending and as such this represents a possible avenue of exploration for future research.

A further possibility is that the binaural beats used in this study were able to elicit a change in the EEG but that this was simply not evident at the temporal (T3/T4) regions examined here. Despite some reports that exposure to binaural beats leads to changes in the temporal regions EEG (Karino et al., 2006; Pratt et al., 2010) others have found changes at both the frontal (Fz: Grose and Mamo, 2011) and central regions (Cz: Hink et al., 1980; Schwarz and Taylor, 2005). Given the disparity in tones it could of course be that distinct frequencies elicit changes in the EEG at different locations. Nevertheless, future research could address this issue by conducting full scalp recordings to monitor for possible changes across different regions.

A final possibility examined here is that the presentation of continuous 1-minute binaural beat tones is either insufficient or incapable of eliciting changes in the EEG. Such an idea is consistent with the findings of Kennerly (1996) who found evidence of entrainment in the EEG occurring on average 5 min after the presentation of the binaural beats. Hence, a continuous tone of 1-minute may be insufficient in length. However, others have shown that prolonged exposure to binaural beats for 20 min and more has also failed to elicit any evidence of a change in the EEG (Goodin et al., 2012, Stevens et al., 2003, Wahbeh et al., 2007). An alternative to the use of continuous tones of varying durations may be to present the binaural beats in short intense bursts. In this way it is the 'intensity' of the tone which facilitates the frequency following the effect in the EEG. It should be noted that this is a speculative proposition as no research has yet directly addressed this issue. Nevertheless, a number of reports have found that short bursts of binaural beats can elicit clear changes in the EEG. For instance, Hink et al. (1980) played tones in short bursts of 100 ms, Schwarz and Taylor (2005) played binaural beats in short bursts of 1200 ms, Pratt et al. (2010) presented binaural beats for bursts of 2000 ms and Grose and Mamo (2011) presented tones in bursts of a little under 2 s. It may be that presenting the binaural beats in short bursts provides a stronger more robust auditory stimulus and as such is more capable of eliciting a response in the EEG. As noted above, this represents a speculative suggestion as no research has directly addressed this question. Nevertheless, future research could address this by directly comparing the effectiveness of the binaural beats delivered using continuous tones vs. short intense bursts.

In conclusion, presentation of interleaved one-minute binaural beats of alpha and beta failed to elicit any clear evidence of a change in the EEG. This severely limits the potential applications of the binaural beats given that a key assumption is that they mediate behavioural change via entrainment of cortical activity. Nevertheless, future research could help to clarify this picture by conducting full scalp recordings and possibly utilising short bursts of binaural beats rather than the continuous and prolonged tones utilised here.

5. Uncited references

Barry et al., 2003
Boutros et al., 1999
Fuentemilla et al., 2006

496 Gruzelier et al., 2010
 497 Pantev et al., 2004
 498 Tanaka et al., 2001
 499 Teplan et al., 2011
 500 Valentino and Dufresne, 1991

501 References

- 502 Anokhin, A., Vogel, F., 1996. EEG alpha rhythm frequency and intelligence in normal
 503 adults. *Intelligence* 23 (1–14).
- 504 Barry, R.J., Cocker, K.I., Anderson, J.W., Gordon, E., Rennie, C., 1992. Does the N100
 505 evoked potential really habituate? Evidence from a paradigm appropriate to a
 506 clinical setting. *International Journal of Psychophysiology* 13 (1), 9–16.
- 507 Barry, R.J., Clarke, A.R., Johnstone, S.J., 2003. A review of electrophysiology in attention-
 508 deficit/hyperactivity disorder: I. Qualitative and quantitative electroencephalography.
 509 *Clinical Neurophysiology* 114 (2), 171–183.
- 510 Bear, D.M., 1979. Temporal lobe epilepsy: a syndrome of sensory–limbic hyperconnection.
 511 *Cortex* 15 (3), 357–384.
- 512 Boutros, N.N., Belger, A., Campbell, D., D'Souza, C., Krystal, J., 1999. Comparison of four
 513 components of sensory gating in schizophrenia and normal subjects: a preliminary
 514 report. *Psychiatry Research* 88, 119–130.
- 515 Brady, B., Stevens, L., 2000. Binaural-beat induced theta EEG activity and hypnotic
 516 susceptibility. *American Journal of Clinical Hypnosis* 43 (1), 53–68.
- 517 Carter, C., 2008. Healthcare performance and the effects of the binaural beats on human
 518 blood pressure and heart rate. *Journal of Hospital Marketing & Public Relations* 18
 519 (2), 213–219.
- 520 David, J.B., Naftali, A., Katz, A., 2010. Tinntrain: a multifactorial treatment for tinnitus
 521 using binaural beats. *The Hearing Journal* 63 (11), 25–28.
- 522 Doppelmayr, M., Klimesch, W., Hodlmoser, K., Sauseng, P., Gruber, W., 2005. Intellig-
 523 ence related upper alpha desynchronisation in a semantic memory task. *Brain*
 524 *Research Bulletin* 66, 171–177.
- 525 Egner, T., Gruzelier, J., 2003. Ecological validity of neurofeedback: modulation of slow
 526 wave EEG enhances musical performance. *NeuroReport* 14 (9), 1221–1224.
- 527 Egner, T., Gruzelier, J., 2004. EEG Biofeedback of low beta band components:
 528 frequency-specific effects on variables of attention and event-related brain poten-
 529 tials. *Clinical Neurophysiology* 115, 131–139.
- 530 Fuchs, T., Birbaumer, N., Lutzenberger, W., Gruzelier, J.H., Kaiser, J., 2003. Neurofeedback
 531 treatment for attention-deficit/hyperactivity disorder in children: a comparison
 532 with methylphenidate. *Applied Psychophysiology and Biofeedback* 28 (1), 1–12.
- 533 Fuentemilla, L.L., Marco-Pallares, J., Grau, C., 2006. Modulation of spectral power and
 534 of phase resetting of EEG contributes differentially to the generation of auditory
 535 event-related potentials. *NeuroImage* 30, 909–916.
- 536 Goodin, P., Ciocriari, J., Baker, K., Carrey, A., Harper, M., Kaufman, J., 2012. A high
 537 density EEG investigation into steady state binaural beat stimulation. *PLoS One* 7
 538 (4), e34789.
- 539 Grose, J.H., Mamo, S.K., 2011. Electrophysiological measurement of binaural beats:
 540 effects of primary tone frequency and observer age. *Ear and Hearing* 32 (5), 1–8.
- 541 Grose, J.H., Buss, E., Hall, J.W., 2012. Binaural beat salience. *Hearing Research* 285 (1–2),
 542 40–45.
- 543 Gruzelier, J., Inoue, A., Smart, R., Steed, A., Steffert, T., 2010. Acting performance and
 544 flow state enhanced with sensory–motor rhythm neurofeedback comparing eco-
 545 logically valid immersive VR and training screen scenarios. *Neuroscience Letters*
 546 480, 112–116.
- 547 Hanslmayr, S., Sauseng, P., Doppelmayr, M., Schabus, M., Klimesch, W., 2005. Increasing
 548 individual upper alpha power by neurofeedback improves cognitive performance
 549 in human subjects. *Applied Psychophysiology and Biofeedback* 30 (1), 1–10.
- 550 Hink, R.F., Koder, K., Yamada, O., Kaga, K., Suzuki, J., 1980. Binaural interaction of a
 551 beating frequency-following response. *Audiology* 19, 36–43.
- 552 Jausovec, N., 1996. Differences in EEG alpha activity related to giftedness. *Intelligence*
 553 23, 159–173.
- 554 Karino, S., Yumoto, M., Itoh, K., Uno, A., Matsuda, M., Yamakawa, K., Sekimoto, M.,
 555 Kaneko, Y., Kaga, K., 2004. Magnetoencephalographic study of human auditory
 556 steady-state responses to binaural beat. *International Congress Series* 1270,
 557 169–172.
- 558 Karino, S., Yumoto, M., Itoh, K., Uno, A., Yamakawa, K., Sekimoto, S., Kagal, K., 2006.
 559 Neuromagnetic responses to binaural beats in human cerebral cortex. *Journal of*
 560 *Neurophysiology* 96, 1927–1938.
- 561 Kennel, S., Taylor, A.G., Lyon, D., Bourguignon, C., 2010. Pilot feasibility study of binaural
 562 auditory beats for reducing symptoms of inattention in children and adolescents
 563 with attention-deficit/hyperactivity disorder. *Journal of Paediatric Nursing* 25
 564 (1), 3–11.
- 565 Kennerly, R., 1996. An empirical investigation into the effect of beta frequency binaural
 566 beat audio signals on four measures of human memory. *Hemi-Synch Journal* 14
 567 (3), 1–4.
- 568 Klimesch, W., Sauseng, P., Gerloff, C., 2003. Enhancing cognitive performance with re-
 569 petitive transcranial magnetic stimulation at human individual alpha frequency.
 570 *European Journal of Neuroscience* 17 (5), 1129–1133.
- 571 Lane, J.D., Kasian, S.J., Owens, J.E., Marsh, G.R., 1998. Binaural auditory beats affect vig-
 572 ilance performance and mood. *Physiology & Behavior* 63 (2), 249–252.
- 573 Lavallee, C.F., Koren, S.A., Persinger, M.A., 2011. A quantitative electroencephalographic
 574 study of meditation and binaural beat entrainment. *Journal of Alternative and*
 575 *Complementary Medicine* 17 (4), 351–355.
- 576 Le Scouarnec, R.P., Poirier, R.M., Owens, J.E., Gauthier, J., Taylor, A.G., Foresman, P.A.,
 577 2001. Use of binaural beat tapes for treatment of anxiety: a pilot study of tape pref-
 578 erence and outcomes. *Alternative Therapies in Health and Medicine* 7 (1), 58–63.
- 579 Lubar, J.F., Swartwood, M.O., Swartwood, J.N., O'Donnell, P.H., 1995a. Evaluation of the
 580 effectiveness of EEG neurofeedback training for ADHD in a clinical setting as mea-
 581 sured by changes in T.O.V.A. scores, behavioural ratings, and WISC-R performance.
 582 *Biofeedback and Self-Regulation* 20 (1), 83–99.
- 583 Lubar, J.F., Swartwood, M.O., Swartwood, J.N., Timmerman, D.L., 1995b. Quantitative EEG
 584 and auditory event related potentials in the evaluation of attention deficit/hyperactivity
 585 disorder: Effects of methylphenidate and implications for neurofeedback training.
 586 *Journal of Psychoeducational Assessment* 143–160.
- 587 Monastra, V.J., Lynn, S.L., Linden, M., Lubar, J.F., Gruzelier, J., La Vaque, T.J., 2006. Electroen-
 588 cephalographic biofeedback in the treatment of attention-deficit/hyperactivity
 589 disorder. *Journal of Neurotherapy* 9 (4), 5–34.
- 590 Moridis, C.N., Klados, M.A., Kokkinakis, I.A., Terzis, V., Economides, A.A., Karlovasitou, A.,
 591 Bamidis, P.D., Karabatakis, V.E., 2010. The impact of audio-visual stimulation on
 592 alpha brain oscillations: An EEG study. *Proceedings of the 10th IEEE International*
 593 *Conference on Information Technology and Applications in Biomedicine*, Corfu,
 594 Greece. 3rd to 5th November.
- 595 Oster, G., 1973. Auditory beats in the brain. *Scientific American* 229 (4), 94–102.
- 596 Padmanabhan, R., Hildreth, A.J., Laws, D., 2005. A prospective, randomised, controlled
 597 study examining binaural beat audio and pre-operative anxiety in patients under-
 598 going general anaesthesia for day case surgery. *Anaesthesia* 60, 874–877.
- 599 Pantev, C., Okamoto, H., Ross, B., Stoll, W., Ciurlia-Guy, E., Kakigi, R., Kubo, T., 2004.
 600 Lateral inhibition and habituation of the human auditory cortex. *European Journal*
 601 *of Neuroscience* 19, 2337–2344.
- 602 Perrott, D.R., Nelson, M.A., 1969. Limits for the detection of binaural beats. *Journal of*
 603 *the Acoustical Society of America* 46, 1477–1481.
- 604 Pratt, H., Starr, A., Michalewski, H.J., Dimitrijevic, A., Bleich, N., Mittelman, N., 2010. A
 605 comparison of auditory evoked potentials to acoustic beats and to binaural beats.
 606 *Hearing Research* 262, 34–44.
- 607 Rasey, H.W., Lubar, J.F., McIntyre, A., Zoffuto, A.C., Abbott, P.L., 1996. EEG Biofeedback
 608 for the enhancement of attentional processing in normal college students. *Journal*
 609 *of Neurotherapy* 1 (3), 15–21.
- 610 Rosenfeld, J.P., Reinhart, A.M., Srivastava, S., 1997. The effects of alpha (10-Hz) and beta
 611 (22 Hz) entrainment stimulation on the alpha and beta EEG bands: individual
 612 differences are critical to prediction of effects. *Applied Psychophysiology and*
 613 *Biofeedback* 22 (1), 3–20.
- 614 Saleh, M., Reimer, J., Penn, R., Ojakangas, C.L., Hatsopoulos, N.G., 2010. Fast and slow os-
 615 cillations in human primary motor cortex predict oncoming behaviourally relevant
 616 cues. *Neuron* 65, 461–471.
- 617 Schwarz, D.W.F., Taylor, P., 2005. Human auditory steady state responses to binaural
 618 and monaural beats. *Clinical Neurophysiology* 116, 658–668.
- 619 Tanaka, H., Monahan, K.D., Seals, D.R., 2001. Age-predicted maximal heart rate
 620 revisited. *Journal of the American College of Cardiology* 37, 153–156.
- 621 Teplan, M., Krakovska, A., Stolz, S., 2006. EEG responses to long-term audio-visual
 622 stimulation. *International Journal of Psychophysiology* 59 (2), 81–90.
- 623 Teplan, M., Krakovska, A., Stolz, S., 2011. Direct effects of audio-visual stimulation on
 624 EEG. *Computer Methods and Programs in Biomedicine* 102, 17–24.
- 625 Timmerman, D.L., Lubar, J.F., Rasey, H.W., Frederick, J.A., 1999. Effects of dominant and
 626 twice-dominant alpha audio-visual stimulation on the cortical EEG. *International*
 627 *Journal of Psychophysiology* 32, 55–61.
- 628 Valentino, D.A., Dufresne, R.L., 1991. Attention tasks and EEG power spectra. *International*
 629 *Journal of Psychophysiology* 11, 299–302.
- 630 Vernon, D., 2009. *Human Potential: Exploring Techniques Used to Enhance Human*
 631 *Performance*. Routledge, London.
- 632 Wahbeh, H., Calabrese, C., Zwicky, H., Zajdel, D., 2007. Binaural beat technology
 633 in humans: a pilot study to assess neuropsychologic, physiologic, and electroen-
 634 cephalographic effects. *Journal of Alternative and Complementary Medicine* 13
 635 (2), 199–206.